



For a MELISA Home Draw In the United States:

First order a test by emailing Denise at denisej@openmedicineinstitute.org to request a MELISA kit and include the following information:

Name:	Email:	Address:
Date of Birth:	Phone:	Ideal days/dates for home draw:

Once you have the kit we arrange for a phlebotomist to draw your blood at home and ship the specimen to Germany. The cost for the home draw and processing is \$150 plus \$76 FedEx to Germany plus shipping kit \$20 for a total \$246. The phlebotomist drops off your specimen at a FedEx location. For home draws outside major metropolitan areas additional mileage cost may apply. We have a network of mobile phlebotomist throughout the United States. This test is drawn only on Monday, Tuesday & Wednesday up until 12 noon. If you have specific questions regarding the test(s) you can email info@melisa.org.

Draw Stations in Northern California:

In California Open Medicine Institute has four draw stations in the Bay Area that have kits and will draw for the MELISA test and they are:

1. San Francisco
2. Lafayette
3. Mountain View
4. Santa Cruz

The cost for the draw and processing at one of these four draw stations is \$50 plus \$76 FedEx to Germany for a total \$126. This test is drawn only on Monday, Tuesday & Wednesday from 8 am until 12 noon.

OMI Draw Station San Francisco

2136 Sutter Street,
San Francisco, CA 94115
(415) 762-1001
8a – 12p

OMI Draw Station Mountain View

2500 Hospital Drive– Building 2,
Mountain View, CA 94040
(650) 691-8633
8a – 12p

OMI Draw Station Lafayette

3190 Old Tunnel Road
Road, Suite B, Lafayette, CA 94549
(925) 290-1822
8a – 12p

OMI Draw Station Santa Cruz (Soquel)

4841 Soquel Drive,
Soquel, CA 95073
Phone: (831) 385-6300
8a – 12p

Payment | Results

10-14 days after the blood has been delivered you will be emailed a PayPal invoice for your MELISA test which can be paid securely online with a credit card (Visa, Mastercard, Amex). Once payment has been made, your results will be emailed to you.

N.B. Patients should not have taken immune suppressant medication within 4 weeks of testing.